

Solving Systems of Linear Equations

MODULE



8



ESSENTIAL QUESTION

How can you use systems of equations to solve real-world problems?



LESSON 8.1

Solving Systems of Linear Equations by Graphing



FL 8.EE.3.8a, 8.EE.3.8c

LESSON 8.2

Solving Systems by Substitution



FL 8.EE.3.8b, 8.EE.3.8c

LESSON 8.3

Solving Systems by Elimination



FL 8.EE.3.8b, 8.EE.3.8c

LESSON 8.4

Solving Systems by Elimination with Multiplication



FL 8.EE.3.8b, 8.EE.3.8c

LESSON 8.5

Solving Special Systems



FL 8.EE.3.8b, 8.EE.3.8c



Real-World Video

The distance contestants in a race travel over time can be modeled by a system of equations. Solving such a system can tell you when one contestant will overtake another who has a head start, as in a boating race or marathon.

my.hrw.com

© Houghton Mifflin Harcourt Publishing Company • Image Credits: © Kenny Ferguson/Alamy Images

GO
DIGITAL
my.hrw.com



my.hrw.com

Go digital with your write-in student edition, accessible on any device.



Math On the Spot

Scan with your smart phone to jump directly to the online edition, video tutor, and more.



Animated Math

Interactively explore key concepts to see how math works.



Personal Math Trainer

Get immediate feedback and help as you work through practice sets.

Are YOU Ready?

Complete these exercises to review skills you will need for this module.



Personal Math Trainer

Online Assessment and Intervention

my.hrw.com

Simplify Algebraic Expressions

EXAMPLE Simplify $5 - 4y + 2x - 6 + y$.

$$\begin{aligned} & -4y + y + 2x - 6 + 5 \\ & -3y + 2x - 1 \end{aligned}$$

Group like terms.
Combine like terms.

Simplify.

1. $14x - 4x + 21$

2. $-y - 4x + 4y$

3. $5.5a - 1 + 21b + 3a$

4. $2y - 3x + 6x - y$

Graph Linear Equations


EXAMPLE Graph $y = -\frac{1}{3}x + 2$.

Step 1: Make a table of values.

x	$y = -\frac{1}{3}x + 2$	(x, y)
0	$y = -\frac{1}{3}(0) + 2 = 2$	(0, 2)
3	$y = -\frac{1}{3}(3) + 2 = 1$	(3, 1)


Step 2: Plot the points.

Step 3: Connect the points with a line.




Graph each equation.


5. $y = 4x - 1$



6. $y = \frac{1}{2}x + 1$




7. $y = -x$



Reading Start-Up

Visualize Vocabulary

Use the ✓ words to complete the graphic.



Understand Vocabulary

Complete the sentences using the preview words.

1. A _____ is any ordered pair that satisfies all the equations in a system.
2. A set of two or more equations that contain two or more variables is called a _____.

Vocabulary

Review Words

- linear equation (*ecuación lineal*)
- ✓ ordered pair (*par ordenado*)
- ✓ slope (*pendiente*)
- slope-intercept form (*forma pendiente intersección*)
- x-axis (*eje x*)
- ✓ x-intercept (*intersección con el eje x*)
- y-axis (*eje y*)
- ✓ y-intercept (*intersección con el eje y*)

Preview Words

- solution of a system of equations (*solución de un sistema de ecuaciones*)
- system of equations (*sistema de ecuaciones*)

Active Reading

Four-Corner Fold Before beginning the module, create a four-corner fold to help you organize what you learn about solving systems of equations. Use the categories "Solving by Graphing," "Solving by Substitution," "Solving by Elimination," and "Solving by Multiplication." As you study this module, note similarities and differences among the four methods. You can use your four-corner fold later to study for tests and complete assignments.





Unpacking the Standards

Understanding the standards and the vocabulary terms in the standards will help you know exactly what you are expected to learn in this module.

FL 8.EE.3.8a

Understand that solutions to a system of two linear equations in two variables correspond to points of intersection of their graphs, because points of intersection satisfy both equations simultaneously.

FL 8.EE.3.8b

Solve systems of two linear equations in two variables algebraically, and estimate solutions by graphing the equations. Solve simple cases by inspection.

Key Vocabulary

solution of a system of equations (*solución de un sistema de ecuaciones*) A set of values that make all equations in a system true.

system of equations (*sistema de ecuaciones*) A set of two or more equations that contain two or more variables.

What It Means to You

You will understand that the points of intersection of two or more graphs represent the solution to a system of linear equations.

UNPACKING EXAMPLE 8.EE.3.8a, 8.EE.3.8b

Use the elimination method.

$$\begin{array}{r} \text{A.} \quad -x = -1 + y \\ \quad x + y = 4 \\ \hline \quad \quad y = y + 3 \end{array}$$

This is never true, so the system has no solution.

The lines never intersect.

$$\begin{array}{r} \text{B.} \quad 2y + x = 1 \\ \quad y - 2 = x \end{array}$$

Use the substitution method.

$$\begin{array}{r} 2y + (y - 2) = 1 \\ 3y - 2 = 1 \\ \quad \quad y = 1 \\ x = y - 2 \\ x = 1 - 2 \\ \quad \quad = -1 \end{array}$$

There is only one solution: $x = -1, y = 2$.

The lines intersect at a single point: $(-1, 2)$.




$$\begin{array}{r} \text{C.} \quad 3y - 6x = 3 \\ \quad y - 2x = 1 \end{array}$$

Use the multiplication method.

$$\begin{array}{r} 3y - 6x = 3 \\ 3y - 6x = 3 \\ \hline \quad \quad 0 = 0 \end{array}$$

This is always true. So the system has infinitely many solutions.

The graphs overlap completely. They are the same line.



Visit my.hrw.com to see all **Florida Math Standards** unpacked.

LESSON 8.1 Solving Systems of Linear Equations by Graphing

 **FL** 8.EE.3.8a

Understand that solutions to a system of two linear equations in two variables correspond to points of intersection of their graphs, because points of intersection satisfy both equations simultaneously. Also 8.EE.3.8, 8.EE.3.8c



ESSENTIAL QUESTION

How can you solve a system of equations by graphing?

EXPLORE ACTIVITY

 **FL** 8.EE.3.8a

Investigating Systems of Equations

You have learned several ways to graph a linear equation in slope-intercept form. For example, you can use the slope and y-intercept or you can find two points that satisfy the equation and connect them with a line.

Slope-intercept form is $y = mx + b$, where m is the slope and b is the y-intercept.

A Graph the pair of equations together: $\begin{cases} y = 3x - 2 \\ y = -2x + 3 \end{cases}$

B Explain how to tell whether $(2, -1)$ is a solution of the equation $y = 3x - 2$ without using the graph.

C Explain how to tell whether $(2, -1)$ is a solution of the equation $y = -2x + 3$ without using the graph.

D Use the graph to explain whether $(2, -1)$ is a solution of each equation.

E Determine if the point of intersection is a solution of both equations.

Point of intersection: (\square, \square)


$$\square = 3\square - 2$$

$$1 = \square$$

$$\square = -2\square + 3$$

$$1 = \square$$

The point of intersection **is / is not** the solution of both equations.





Math On the Spot

my.hrw.com

Solving Systems Graphically

An ordered pair (x, y) is a solution of an equation in two variables if substituting the x - and y -values into the equation results in a true statement. A **system of equations** is a set of equations that have the same variables. An ordered pair is a **solution of a system of equations** if it is a solution of every equation in the set.

Since the graph of an equation represents all ordered pairs that are solutions of the equation, if a point lies on the graphs of two equations, the point is a solution of both equations and is, therefore, a solution of the system.

EXAMPLE 1



FL 8.EE.3.8

Solve each system by graphing.


A $\begin{cases} y = -x + 4 \\ y = 3x \end{cases}$

STEP 1 Start by graphing each equation.

STEP 2 Find the point of intersection of the two lines. It appears to be $(1, 3)$. Check by substitution to determine if it is a solution to both equations.

$$\begin{array}{ll} y = -x + 4 & y = 3x \\ 3 \stackrel{?}{=} -(1) + 4 & 3 \stackrel{?}{=} 3(1) \\ 3 = 3 \checkmark & 3 = 3 \checkmark \end{array}$$

The solution of the system is $(1, 3)$.




B $\begin{cases} y = 3x - 3 \\ y = 3(x - 1) \end{cases}$

STEP 1 Start by graphing each equation.

STEP 2 Identify any ordered pairs that are solutions of both equations.

The graphs of the equations are the same line. So, every ordered pair that is a solution of one equation is also a solution of the other equation. The system has infinitely many solutions.



My Notes

Reflect

1. A system of linear equations has infinitely many solutions. Does that mean any ordered pair in the coordinate plane is a solution?


2. Can you show algebraically that both equations in part B represent the same line? If so, explain how.

YOUR TURN

Solve each system by graphing. Check by substitution.


3. $\begin{cases} y = -x + 2 \\ y = -4x - 1 \end{cases}$ _____

Check:



4. $\begin{cases} y = -2x + 5 \\ y = 3x \end{cases}$ _____

Check:





Math On the Spot

my.hrw.com

Solving Problems Using Systems of Equations

When using graphs to solve a system of equations, it is best to rewrite both equations in slope-intercept form for ease of graphing.

To write an equation in slope-intercept form starting from $ax + by = c$:

$$ax + by = c$$

$$by = c - ax \quad \text{Subtract } ax \text{ from both sides.}$$

$$y = \frac{c}{b} - \frac{ax}{b} \quad \text{Divide both sides by } b.$$

$$y = -\frac{a}{b}x + \frac{c}{b} \quad \text{Rearrange the equation.}$$

EXAMPLE 2



FL 8.EE.3.8c, 8.EE.3.8

Keisha and her friends visit the concession stand at a football game. The stand charges \$2 for a hot dog and \$1 for a drink. The friends buy a total of 8 items for \$11. Tell how many hot dogs and how many drinks they bought.

STEP 1

Let x represent the number of hot dogs they bought and let y represent the number of drinks they bought.

Write an equation representing the **number of items they purchased**.

$$\text{Number of hot dogs} + \text{Number of drinks} = \text{Total items}$$

$$x + y = 8$$

Write an equation representing the **money spent on the items**.

$$\text{Cost of 1 hot dog times number of hot dogs} + \text{Cost of 1 drink times number of drinks} = \text{Total cost}$$

$$2x + 1y = 11$$

STEP 2

Write the equations in slope-intercept form. Then graph.

$$x + y = 8$$

$$y = 8 - x$$


$$y = -x + 8$$

$$2x + 1y = 11$$

$$1y = 11 - 2x$$

$$y = -2x + 11$$

Graph the equations $y = -x + 8$ and $y = -2x + 11$.



© Houghton Mifflin Harcourt Publishing Company • Image Credits:
© Veronica Ankorhorn/Houghton Mifflin Harcourt

STEP 3

Use the graph to identify the solution of the system of equations. Check your answer by substituting the ordered pair into both equations.

Apparent solution: (3, 5)

Check:

$$\begin{array}{rcl} x + y = 8 & & 2x + y = 11 \\ 3 + 5 \stackrel{?}{=} 8 & & 2(3) + 5 \stackrel{?}{=} 11 \\ 8 = 8 \checkmark & & 11 = 11 \checkmark \end{array}$$

The point (3, 5) is a solution of both equations.

STEP 4

Interpret the solution in the original context.

- Keisha and her friends bought 3 hot dogs and 5 drinks.




Reflect

5. **Conjecture** Why do you think the graph is limited to the first quadrant?

YOUR TURN

6. During school vacation, Marquis wants to go bowling and to play laser tag. He wants to play 6 total games but needs to figure out how many of each he can play if he spends exactly \$20. Each game of bowling is \$2 and each game of laser tag is \$4.
- a. Let x represent the number of games Marquis bowls and let y represent the number of games of laser tag Marquis plays. Write a system of equations that describes the situation. Then write the equations in slope-intercept form.


- b. Graph the solutions of both equations.
- c. How many games of bowling and how many games of laser tag will Marquis play?




Guided Practice

Solve each system by graphing. (Example 1)

1. $\begin{cases} y = 3x - 4 \\ y = x + 2 \end{cases}$ _____



2. $\begin{cases} x - 3y = 2 \\ -3x + 9y = -6 \end{cases}$ _____




3. Mrs. Morales wrote a test with 15 questions covering spelling and vocabulary. Spelling questions (x) are worth 5 points and vocabulary questions (y) are worth 10 points. The maximum number of points possible on the test is 100. (Example 2)

a. Write an equation in slope-intercept form to represent the number of questions on the test.

b. Write an equation in slope-intercept form to represent the total number of points on the test.

c. Graph the solutions of both equations.

d. Use your graph to tell how many of each question type are on the test.




ESSENTIAL QUESTION CHECK-IN

4. When you graph a system of linear equations, why does the intersection of the two lines represent the solution of the system?

8.1 Independent Practice



FL 8.EE.3.8, 8.EE.3.8a, 8.EE.3.8c



Personal Math Trainer

Online Assessment and Intervention

my.hrw.com


5. Vocabulary A _____ is a set of equations that have the same variables.

6. Eight friends started a business. They will wear either a baseball cap or a shirt imprinted with their logo while working. They want to spend exactly \$36 on the shirts and caps. Shirts cost \$6 each and caps cost \$3 each.

a. Write a system of equations to describe the situation. Let x represent the number of shirts and let y represent the number of caps.

b. Graph the system. What is the solution and what does it represent?

Business Logo Wear




7. Multistep The table shows the cost for bowling at two bowling alleys.

	Shoe Rental Fee	Cost per Game
Bowl-o-Rama	\$2.00	\$2.50
Bowling Pinz	\$4.00	\$2.00

a. Write a system of equations, with one equation describing the cost to bowl at Bowl-o-Rama and the other describing the cost to bowl at Bowling Pinz. For each equation, let x represent the number of games played and let y represent the total cost.

b. Graph the system. What is the solution and what does it represent?

Cost of Bowling



8. **Multi-Step** Jeremy runs 7 miles per week and increases his distance by 1 mile each week. Tony runs 3 miles per week and increases his distance by 2 miles each week. In how many weeks will Jeremy and Tony be running the same distance? What will that distance be?

9. **Critical Thinking** Write a real-world situation that could be represented by the system of equations shown below.

$$\begin{cases} y = 4x + 10 \\ y = 3x + 15 \end{cases}$$



FOCUS ON HIGHER ORDER THINKING

10. **Multistep** The table shows two options provided by a high-speed Internet provider.

	Setup Fee (\$)	Cost per Month (\$)
Option 1	50	30
Option 2	No setup fee	\$40

- a. In how many months will the total cost of both options be the same? What will that cost be?
- b. If you plan to cancel your Internet service after 9 months, which is the cheaper option? Explain.

11. **Draw Conclusions** How many solutions does the system formed by $x - y = 3$ and $ay - ax + 3a = 0$ have for a nonzero number a ? Explain.

Work Area

LESSON 8.2 Solving Systems by Substitution

 **FL** 8.EE.3.8b

Solve systems of two linear equations in two variables algebraically, and estimate solutions by graphing the equations. ... Also 8.EE.3.8c



ESSENTIAL QUESTION

How do you use substitution to solve a system of linear equations?

Solving a Linear System by Substitution

The **substitution method** is used to solve systems of linear equations by solving an equation for one variable and then substituting the resulting expression for that variable into the other equation. The steps for this method are as follows:

1. Solve one of the equations for one of its variables.
2. Substitute the expression from step 1 into the other equation and solve for the other variable.
3. Substitute the value from step 2 into either original equation and solve to find the value of the variable in step 1.

EXAMPLE 1

 **FL** 8.EE.3.8b

Solve the system of linear equations by substitution. Check your answer.

$$\begin{cases} -3x + y = 1 \\ 4x + y = 8 \end{cases}$$

STEP 1 Solve an equation for one variable.

$$-3x + y = 1 \quad \text{Select one of the equations.}$$

$$y = 3x + 1 \quad \text{Solve for the variable } y. \text{ Isolate } y \text{ on one side.}$$

STEP 2 Substitute the expression for y in the other equation and solve.

$$4x + (3x + 1) = 8 \quad \text{Substitute the expression for the variable } y.$$

$$7x + 1 = 8 \quad \text{Combine like terms.}$$

$$7x = 7 \quad \text{Subtract 1 from each side.}$$

$$x = 1 \quad \text{Divide each side by 7.}$$

STEP 3 Substitute the value of x you found into one of the equations and solve for the other variable, y .

$$-3(1) + y = 1 \quad \text{Substitute the value of } x \text{ into the first equation.}$$

$$-3 + y = 1 \quad \text{Simplify.}$$

$$y = 4 \quad \text{Add 3 to each side.}$$

So, $(1, 4)$ is the solution of the system.



My Notes

STEP 4 Check the solution by graphing.

$$-3x + y = 1$$


$$4x + y = 8$$

$$x\text{-intercept: } -\frac{1}{3}$$

$$x\text{-intercept: } 2$$

$$y\text{-intercept: } 1$$

$$y\text{-intercept: } 8$$



The point of intersection is (1, 4).

Reflect

1. Is it more efficient to solve $-3x + y = 1$ for x ? Why or why not?

2. Is there another way to solve the system?

3. What is another way to check your solution?

YOUR TURN

Solve each system of linear equations by substitution.

4.
$$\begin{cases} 3x + y = 11 \\ -2x + y = 1 \end{cases}$$

5.
$$\begin{cases} 2x - 3y = -24 \\ x + 6y = 18 \end{cases}$$

6.
$$\begin{cases} x - 2y = 5 \\ 3x - 5y = 8 \end{cases}$$



Personal Math Trainer

Online Assessment and Intervention

my.hrw.com

Using a Graph to Estimate the Solution of a System

You can use a graph to estimate the solution of a system of equations before solving the system algebraically.




EXAMPLE 2



FL 8.EE.3.8b

Solve the system $\begin{cases} x - 4y = 4 \\ 2x - 3y = -3 \end{cases}$.

STEP 1 Sketch a graph of each equation by substituting values for x and generating values of y .



STEP 2 Find the intersection of the lines. The lines appear to intersect near $(-5, -2)$.

STEP 3 Solve the system algebraically.

Solve $x - 4y = 4$ for x .	Substitute to find y .	Substitute to find x .
$x - 4y = 4$	$2(4 + 4y) - 3y = -3$	$x = 4 + 4y$
$x = 4 + 4y$	$8 + 8y - 3y = -3$	$= 4 + 4\left(-\frac{11}{5}\right)$
	$8 + 5y = -3$	$= \frac{20 - 44}{5}$
	$5y = -11$	$= -\frac{24}{5}$
	$y = -\frac{11}{5}$	

The solution is $\left(-\frac{24}{5}, -\frac{11}{5}\right)$.

STEP 4 Use the estimate you made using the graph to judge the reasonableness of your solution.

$-\frac{24}{5}$ is close to the estimate of -5 , and $-\frac{11}{5}$ is close to the estimate of -2 , so the solution seems reasonable.




Math Talk

Mathematical Practices

In Step 2, how can you tell that $(-5, -2)$ is not the solution?

YOUR TURN

7. Estimate the solution of the system $\begin{cases} x + y = 4 \\ 2x - y = 6 \end{cases}$ by sketching a graph of each linear function. Then solve the system algebraically. Use your estimate to judge the reasonableness of your solution.



The estimated solution is _____.

The algebraic solution is _____.

The solution is/is not reasonable because

Solving Problems with Systems of Equations




EXAMPLE 3



FL 8.EE.3.8c

As part of Class Day, the eighth grade is doing a treasure hunt. Each team is given the following riddle and map. At what point is the treasure located?

There's pirate treasure to be found. So search on the island, all around. Draw a line through A and B. Then another through C and D. Dance a jig, "X" marks the spot. Where the lines intersect, that's the treasure's plot!



STEP 1

Give the coordinates of each point and find the slope of the line through each pair of points.

A: (-2, -1) C: (-1, 4)

B: (2, 5) D: (1, -4)

Slope:	Slope:
$\frac{5 - (-1)}{2 - (-2)} = \frac{6}{4}$	$\frac{-4 - 4}{1 - (-1)} = \frac{-8}{2}$
$= \frac{3}{2}$	$= -4$

Math Talk

Mathematical Practices

Where do the lines appear to intersect? How is this related to the solution?

STEP 2 Write equations in slope-intercept form describing the line through points A and B and the line through points C and D .

Line through A and B :

Use the slope and a point to find b .

$$5 = \left(\frac{3}{2}\right)2 + b$$

$$b = 2$$

The equation is $y = \frac{3}{2}x + 2$.

Line through C and D :

Use the slope and a point to find b .

$$4 = -4(-1) + b$$

$$b = 0$$

The equation is $y = -4x$.

STEP 3 Solve the system algebraically.

Substitute $\frac{3}{2}x + 2$ for y in $y = -4x$ to find x .

$$\frac{3}{2}x + 2 = -4x$$

$$\frac{11}{2}x = -2$$

$$x = -\frac{4}{11}$$

Substitute to find x .

$$y = -4\left(-\frac{4}{11}\right) = \frac{16}{11}$$

The solution is $\left(-\frac{4}{11}, \frac{16}{11}\right)$.

YOUR TURN

8. Ace Car Rental rents cars for x dollars per day plus y dollars for each mile driven. Carlos rented a car for 4 days, drove it 160 miles, and spent \$120. Vanessa rented a car for 1 day, drove it 240 miles, and spent \$80. Write equations to represent Carlos's expenses and Vanessa's expenses. Then solve the system and tell what each number represents.



**Personal
Math Trainer**

Online Assessment
and Intervention

my.hrw.com

Guided Practice

Solve each system of linear equations by substitution. (Example 1)

1. $\begin{cases} 3x - 2y = 9 \\ y = 2x - 7 \end{cases}$ _____

2. $\begin{cases} y = x - 4 \\ 2x + y = 5 \end{cases}$ _____

3. $\begin{cases} x + 4y = 6 \\ y = -x + 3 \end{cases}$ _____

4. $\begin{cases} x + 2y = 6 \\ x - y = 3 \end{cases}$ _____

Solve each system. Estimate the solution first. (Example 2)

5. $\begin{cases} 6x + y = 4 \\ x - 4y = 19 \end{cases}$

Estimate _____

Solution _____

6. $\begin{cases} x + 2y = 8 \\ 3x + 2y = 6 \end{cases}$

Estimate _____

Solution _____

7. $\begin{cases} 3x + y = 4 \\ 5x - y = 22 \end{cases}$

Estimate _____

Solution _____

8. $\begin{cases} 2x + 7y = 2 \\ x + y = -1 \end{cases}$

Estimate _____

Solution _____

9. Adult tickets to Space City amusement park cost x dollars. Children's tickets cost y dollars. The Henson family bought 3 adult and 1 child tickets for \$163. The Garcia family bought 2 adult and 3 child tickets for \$174. (Example 3)

- a. Write equations to represent the Hensons' cost and the Garcias' cost.

Hensons' cost: _____

Garcias' cost: _____

- b. Solve the system.

adult ticket price: _____

child ticket price: _____




ESSENTIAL QUESTION CHECK-IN

10. How can you decide which variable to solve for first when you are solving a linear system by substitution?

8.2 Independent Practice



FL 8.EE.3.8b, 8.EE.3.8c




Personal Math Trainer

Online Assessment and Intervention

my.hrw.com

- 11. Check for Reasonableness** Zach solves the system $\begin{cases} x + y = -3 \\ x - y = 1 \end{cases}$ and finds the solution $(1, -2)$. Use a graph to explain whether Zach's solution is reasonable.



- 12. Represent Real-World Problems** Angelo bought apples and bananas at the fruit stand. He bought 20 pieces of fruit and spent \$11.50. Apples cost \$0.50 and bananas cost \$0.75 each.

- a.** Write a system of equations to model the problem. (Hint: One equation will represent the number of pieces of fruit. A second equation will represent the money spent on the fruit.)

- b.** Solve the system algebraically. Tell how many apples and bananas Angelo bought.



- 13. Represent Real-World Problems** A jar contains n nickels and d dimes. There is a total of 200 coins in the jar. The value of the coins is \$14.00. How many nickels and how many dimes are in the jar?

- 14. Multistep** The graph shows a triangle formed by the x -axis, the line $3x - 2y = 0$, and the line $x + 2y = 10$. Follow these steps to find the area of the triangle.

- a.** Find the coordinates of point A by solving the system $\begin{cases} 3x - 2y = 0 \\ x + 2y = 10 \end{cases}$.

Point A : _____


- b.** Use the coordinates of point A to find the height of the triangle.

height: _____

- c.** What is the length of the base of the triangle?

base: _____

- d.** What is the area of the triangle? _____



15. Jed is graphing the design for a kite on a coordinate grid. The four vertices of the kite are at $A\left(-\frac{4}{3}, \frac{2}{3}\right)$, $B\left(\frac{14}{3}, -\frac{4}{3}\right)$, $C\left(\frac{14}{3}, -\frac{16}{3}\right)$, and $D\left(\frac{2}{3}, -\frac{16}{3}\right)$. One kite strut will connect points A and C . The other will connect points B and D . Find the point where the struts cross.



H.O.T. FOCUS ON HIGHER ORDER THINKING

16. **Analyze Relationships** Consider the system $\begin{cases} 6x - 3y = 15 \\ x + 3y = -8 \end{cases}$. Describe three different substitution methods that can be used to solve this system. Then solve the system.

17. **Communicate Mathematical Ideas** Explain the advantages, if any, that solving a system of linear equations by substitution has over solving the same system by graphing.

18. **Persevere in Problem Solving** Create a system of equations of the form $\begin{cases} Ax + By = C \\ Dx + Ey = F \end{cases}$ that has $(7, -2)$ as its solution. Explain how you found the system.

Work Area

LESSON 8.3 Solving Systems by Elimination

 **FL** 8.EE.3.8b

Solve systems of two linear equations in two variables algebraically, . . . Also 8.EE.3.8c



ESSENTIAL QUESTION

How do you solve a system of linear equations by adding or subtracting?

Solving a Linear System by Adding

The **elimination method** is another method used to solve a system of linear equations. In this method, one variable is *eliminated* by adding or subtracting the two equations of the system to obtain a single equation in one variable. The steps for this method are as follows:

1. Add or subtract the equations to eliminate one variable.
2. Solve the resulting equation for the other variable.
3. Substitute the value into either original equation to find the value of the eliminated variable.



EXAMPLE 1

 **FL** 8.EE.3.8b

Solve the system of equations by adding. Check your answer.

$$\begin{cases} 2x - 3y = 12 \\ x + 3y = 6 \end{cases}$$

STEP 1 Add the equations.

$$\begin{array}{r} 2x - 3y = 12 \\ + x + 3y = 6 \\ \hline 3x + 0 = 18 \\ 3x = 18 \\ \frac{3x}{3} = \frac{18}{3} \\ x = 6 \end{array}$$

Write the equations so that like terms are aligned.
Notice that the terms $-3y$ and $3y$ are opposites.
Add to eliminate the variable y .
Simplify and solve for x .
Divide each side by 3.
Simplify.

STEP 2 Substitute the solution into one of the original equations and solve for y .

$$\begin{array}{r} x + 3y = 6 \\ 6 + 3y = 6 \\ 3y = 0 \\ y = 0 \end{array}$$

Use the second equation.
Substitute 6 for the variable x .
Subtract 6 from each side.
Divide each side by 3 and simplify.

My Notes

STEP 3

Write the solution as an ordered pair: (6, 0)

STEP 4

Check the solution by graphing.

$$2x - 3y = 12$$


$$x + 3y = 6$$

x-intercept: 6

x-intercept: 6

y-intercept: -4

y-intercept: 2



The point of intersection is (6, 0).

Math Talk
Mathematical Practices

Is it better to check a solution by graphing or by substituting the values in the original equations?

Reflect

- Can this linear system be solved by subtracting one of the original equations from the other? Why or why not?

- What is another way to check your solution?

YOUR TURN

Solve each system of equations by adding. Check your answers.

3. $\begin{cases} x + y = -1 \\ x - y = 7 \end{cases}$

4. $\begin{cases} 2x + 2y = -2 \\ 3x - 2y = 12 \end{cases}$

5. $\begin{cases} 6x + 5y = 4 \\ -6x + 7y = 20 \end{cases}$

Personal Math Trainer
Online Assessment and Intervention
my.hrw.com

Solving a Linear System by Subtracting

If both equations contain the same x - or y -term, you can solve by subtracting.



Math On the Spot

my.hrw.com

EXAMPLE 2



FL 8.EE.3.8b

Solve the system of equations by subtracting. Check your answer.

$$\begin{cases} 3x + 3y = 6 \\ 3x - y = -6 \end{cases}$$

STEP 1 Subtract the equations.

$$\begin{array}{r} 3x + 3y = 6 \\ -(3x - y = -6) \\ \hline 0 + 4y = 12 \\ 4y = 12 \\ y = 3 \end{array}$$

Write the equations so that like terms are aligned.
Notice that both equations contain the term $3x$.
Subtract to eliminate the variable x .
Simplify and solve for y .
Divide each side by 4 and simplify.

STEP 2 Substitute the solution into one of the original equations and solve for x .

$$\begin{array}{r} 3x - y = -6 \\ 3x - 3 = -6 \\ 3x = -3 \\ x = -1 \end{array}$$


Use the second equation.
Substitute 3 for the variable y .
Add 3 to each side.
Divide each side by 3 and simplify.

STEP 3 Write the solution as an ordered pair: $(-1, 3)$

STEP 4 Check the solution by graphing.

$$\begin{array}{ll} 3x + 3y = 6 & 3x - y = -6 \\ x\text{-intercept: } 2 & x\text{-intercept: } -2 \\ y\text{-intercept: } 2 & y\text{-intercept: } 6 \end{array}$$

The point of intersection is $(-1, 3)$.




Reflect

6. What would happen if you added the original equations?

My Notes

7. How can you decide whether to add or subtract to eliminate a variable in a linear system? Explain your reasoning.



Personal Math Trainer
Online Assessment and Intervention
my.hrw.com

YOUR TURN

Solve each system of equations by subtracting. Check your answers.

8. $\begin{cases} 6x - 3y = 6 \\ 6x + 8y = -16 \end{cases}$

9. $\begin{cases} 4x + 3y = 19 \\ 6x + 3y = 33 \end{cases}$

10. $\begin{cases} 2x + 6y = 17 \\ 2x - 10y = 9 \end{cases}$



Math On the Spot
my.hrw.com

Solving Problems with Systems of Equations

Many real-world situations can be modeled and solved with a system of equations.

EXAMPLE 3



FL

8.EE.3.8c

The Polar Bear Club wants to buy snowshoes and camp stoves. The club will spend \$554.50 to buy them at Top Sports and \$602.00 to buy them at Outdoor Explorer, before taxes, but Top Sports is farther away. How many of each item does the club intend to buy?



	Snowshoes	Camp Stoves
Top Sports	\$79.50 per pair	\$39.25
Outdoor Explorer	\$89.00 per pair	\$39.25

STEP 1 Choose variables and write a system of equations.
 Let x represent the number of pairs of snowshoes.
 Let y represent the number of camp stoves.

$$\text{Top Sports cost: } 79.50x + 39.25y = 554.50$$

$$\text{Outdoor Explorer cost: } 89.00x + 39.25y = 602.00$$

STEP 2 Subtract the equations.

$$79.50x + 39.25y = 554.50$$

$$-(89.00x + 39.25y = 602.00)$$

$$\hline -9.50x + 0 = -47.50$$

$$-9.50x = -47.50$$

$$\frac{-9.50x}{-9.50} = \frac{-47.50}{-9.50}$$

$$x = 5$$

Both equations contain the term $39.25y$.

Subtract to eliminate the variable y .

Simplify and solve for x .

Divide each side by -9.50 .

Simplify.

STEP 3 Substitute the solution into one of the original equations and solve for y .

$$79.50x + 39.25y = 554.50$$

$$79.50(5) + 39.25y = 554.50$$

$$397.50 + 39.25y = 554.50$$

$$39.25y = 157.00$$

$$\frac{39.25y}{39.25} = \frac{157.00}{39.25}$$

$$y = 4$$

Use the first equation.

Substitute 5 for the variable x .

Multiply.

Subtract 397.50 from each side.

Divide each side by 39.25.

Simplify.

STEP 4 Write the solution as an ordered pair: (5, 4)

The club intends to buy 5 pairs of snowshoes and 4 camp stoves.

YOUR TURN

11. At the county fair, the Baxter family bought 6 hot dogs and 4 juice drinks for \$16.70. The Farley family bought 3 hot dogs and 4 juice drinks for \$10.85. Find the price of a hot dog and the price of a juice drink.



Guided Practice

1. Solve the system $\begin{cases} 4x + 3y = 1 \\ x - 3y = -11 \end{cases}$ by adding. (Example 1)

STEP 1 Add the equations.

$$\begin{array}{r} 4x + 3y = 1 \\ + \quad x - 3y = -11 \\ \hline \end{array}$$

Write the equations so that like terms are aligned.

$$5x + \boxed{} = \boxed{}$$

Add to eliminate the variable $\boxed{}$.

$$5x = \boxed{}$$

Simplify and solve for x .

$$x = \boxed{}$$

Divide both sides by $\boxed{}$ and simplify.

STEP 2 Substitute into one of the original equations and solve for y .

$y = \boxed{}$ So, $\boxed{}$ is the solution of the system.

Solve each system of equations by adding or subtracting. (Examples 1, 2)

2. $\begin{cases} x + 2y = -2 \\ -3x + 2y = -10 \end{cases}$

3. $\begin{cases} 3x + y = 23 \\ 3x - 2y = 8 \end{cases}$

4. $\begin{cases} -4x - 5y = 7 \\ 3x + 5y = -14 \end{cases}$

5. $\begin{cases} x - 2y = -19 \\ 5x + 2y = 1 \end{cases}$

6. $\begin{cases} 3x + 4y = 18 \\ -2x + 4y = 8 \end{cases}$

7. $\begin{cases} -5x + 7y = 11 \\ -5x + 3y = 19 \end{cases}$

8. The Green River Freeway has a minimum and a maximum speed limit. Tony drove for 2 hours at the minimum speed limit and 3.5 hours at the maximum limit, a distance of 355 miles. Rae drove 2 hours at the minimum speed limit and 3 hours at the maximum limit, a distance of 320 miles. What are the two speed limits? (Example 3)

- a. Write equations to represent Tony's distance and Rae's distance.

Tony: _____

Rae: _____

- b. Solve the system.

minimum speed limit: _____ maximum speed limit: _____




ESSENTIAL QUESTION CHECK-IN

9. Can you use addition or subtraction to solve any system? Explain.


8.3 Independent Practice



FL 8.EE.3.8b, 8.EE.3.8c



- 10. Represent Real-World Problems** Marta bought new fish for her home aquarium. She bought 3 guppies and 2 platies for a total of \$13.95. Hank also bought guppies and platies for his aquarium. He bought 3 guppies and 4 platies for a total of \$18.33. Find the price of a guppy and the price of a platy.



- 11. Represent Real-World Problems** The rule for the number of fish in a home aquarium is 1 gallon of water for each inch of fish length. Marta's aquarium holds 13 gallons and Hank's aquarium holds 17 gallons. Based on the number of fish they bought in Exercise 10, how long is a guppy and how long is a platy?



- 12.** Line m passes through the points $(6, 1)$ and $(2, -3)$. Line n passes through the points $(2, 3)$ and $(5, -6)$. Find the point of intersection of these lines.

- 13. Represent Real-World Problems** Two cars got an oil change at the same auto shop. The shop charges customers for each quart of oil plus a flat fee for labor. The oil change for one car required 5 quarts of oil and cost \$22.45. The oil change for the other car required 7 quarts of oil and cost \$25.45. How much is the labor fee and how much is each quart of oil?

- 14. Represent Real-World Problems** A sales manager noticed that the number of units sold for two T-shirt styles, style A and style B, was the same during June and July. In June, total sales were \$2779 for the two styles, with A selling for \$15.95 per shirt and B selling for \$22.95 per shirt. In July, total sales for the two styles were \$2385.10, with A selling at the same price and B selling at a discount of 22% off the June price. How many T-shirts of each style were sold in June and July combined?

- 15. Represent Real-World Problems** Adult tickets to a basketball game cost \$5. Student tickets cost \$1. A total of \$2,874 was collected on the sale of 1,246 tickets. How many of each type of ticket were sold?





16. **Communicate Mathematical Ideas** Is it possible to solve the system $\begin{cases} 3x - 2y = 10 \\ x + 2y = 6 \end{cases}$ by using substitution? If so, explain how. Which method, substitution or elimination, is more efficient? Why?

17. Jenny used substitution to solve the system $\begin{cases} 2x + y = 8 \\ x - y = 1 \end{cases}$. Her solution is shown below.

- Step 1** $y = -2x + 8$ Solve the first equation for y .
- Step 2** $2x + (-2x + 8) = 8$ Substitute the value of y in an original equation.
- Step 3** $2x - 2x + 8 = 8$ Use the Distributive Property.
- Step 4** $8 = 8$ Simplify.

- a. **Explain the Error** Explain the error Jenny made. Describe how to correct it.

- b. **Communicate Mathematical Ideas** Would adding the equations have been a better method for solving the system? If so, explain why.

LESSON 8.4 Solving Systems by Elimination with Multiplication

 **FL** 8.EE.3.8b

Solve systems of two linear equations in two variables algebraically, ... Also 8.EE.3.8c



ESSENTIAL QUESTION

How do you solve a system of linear equations by multiplying?

Solving a System by Multiplying and Adding

In some linear systems, neither variable can be eliminated by adding or subtracting the equations directly. In systems like these, you need to multiply one of the equations by a constant so that adding or subtracting the equations will eliminate one variable. The steps for this method are as follows:

1. Decide which variable to eliminate.
2. Multiply one equation by a constant so that adding or subtracting will eliminate that variable.
3. Solve the system using the elimination method.



EXAMPLE 1

 **FL** 8.EE.3.8b

Solve the system of equations by multiplying and adding.

$$\begin{cases} 2x + 10y = 2 \\ 3x - 5y = -17 \end{cases}$$

STEP 1

The coefficient of y in the first equation, 10, is 2 times the coefficient of y , 5, in the second equation. Also, the y -term in the first equation is being added, while the y -term in the second equation is being subtracted. To eliminate the y -terms, multiply the second equation by 2 and add this new equation to the first equation.

$$2(3x - 5y = -17)$$

Multiply each term in the second equation by 2 to get opposite coefficients for the y -terms.

$$6x - 10y = -34$$

Simplify.

$$6x - 10y = -34$$

$$+ 2x + 10y = 2$$

Add the first equation to the new equation.

$$8x + 0y = -32$$

Add to eliminate the variable y .

$$8x = -32$$

Simplify and solve for x .

$$\frac{8x}{8} = \frac{-32}{8}$$

Divide each side by 8.

$$x = -4$$

Simplify.

My Notes

STEP 2

Substitute the solution into one of the original equations and solve for y .

$$2x + 10y = 2 \quad \text{Use the first equation.}$$

$$2(-4) + 10y = 2 \quad \text{Substitute } -4 \text{ for the variable } x.$$

$$-8 + 10y = 2 \quad \text{Simplify.}$$

$$10y = 10 \quad \text{Add 8 to each side.}$$

$$y = 1 \quad \text{Divide each side by 10 and simplify.}$$

STEP 3

Write the solution as an ordered pair: $(-4, 1)$

STEP 4

Check your answer algebraically.

Substitute -4 for x and 1 for y in the original system.

$$\begin{cases} 2x + 10y = 2 \rightarrow 2(-4) + 10(1) = -8 + 10 = 2 \checkmark \\ 3x - 5y = -17 \rightarrow 3(-4) - 5(1) = -12 - 5 = -17 \checkmark \end{cases}$$

The solution is correct.

Math Talk**Mathematical Practices**

When you check your answer algebraically, why do you substitute your values for x and y into the original system? Explain.

Reflect

- How can you solve this linear system by subtracting? Which is more efficient, adding or subtracting? Explain your reasoning.

- Can this linear system be solved by adding or subtracting without multiplying? Why or why not?

- What would you need to multiply the second equation by to eliminate x by adding? Why might you choose to eliminate y instead of x ?

YOUR TURN

Solve each system of equations by multiplying and adding.

4. $\begin{cases} 5x + 2y = -10 \\ 3x + 6y = 66 \end{cases}$

5. $\begin{cases} 4x + 2y = 6 \\ 3x - y = -8 \end{cases}$

6. $\begin{cases} -6x + 9y = -12 \\ 2x + y = 0 \end{cases}$



Personal Math Trainer

Online Assessment and Intervention

my.hrw.com

Solving a System by Multiplying and Subtracting

You can solve some systems of equations by multiplying one equation by a constant and then subtracting.



EXAMPLE 2

FL 8.EE.3.8b

Solve the system of equations by multiplying and subtracting.

$$\begin{cases} 6x + 5y = 7 \\ 2x - 4y = -26 \end{cases}$$

STEP 1 Multiply the second equation by 3 and subtract this new equation from the first equation.

$$3(2x - 4y) = -26$$

Multiply each term in the second equation by 3 to get the same coefficients for the x-terms.

$$6x - 12y = -78$$

Simplify.

$$\begin{array}{r} 6x + 5y = 7 \\ -(6x - 12y = -78) \\ \hline \end{array}$$

Subtract the new equation from the first equation.

$$0x + 17y = 85$$

Subtract to eliminate the variable x.

$$17y = 85$$

Simplify and solve for y.

$$\frac{17y}{17} = \frac{85}{17}$$

Divide each side by 17.

$$y = 5$$

Simplify.

STEP 2 Substitute the solution into one of the original equations and solve for x.

$$6x + 5y = 7$$

Use the first equation.

$$6x + 5(5) = 7$$

Substitute 5 for the variable y.

$$6x + 25 = 7$$

Simplify.

$$6x = -18$$

Subtract 25 from each side.

$$x = -3$$

Divide each side by 6 and simplify.


STEP 3 Write the solution as an ordered pair: $(-3, 5)$

STEP 4 Check your answer algebraically.
Substitute -3 for x and 5 for y in the original system.

$$\begin{cases} 6x + 5y = 7 \rightarrow 6(-3) + 5(5) = -18 + 25 = 7 \checkmark \\ 2x - 4y = -26 \rightarrow 2(-3) - 4(5) = -6 - 20 = -26 \checkmark \end{cases}$$

The solution is correct.

My Notes



Personal Math Trainer
Online Assessment and Intervention
my.hrw.com

YOUR TURN

Solve each system of equations by multiplying and subtracting.

7. $\begin{cases} 3x - 7y = 2 \\ 6x - 9y = 9 \end{cases}$

8. $\begin{cases} -3x + y = 11 \\ 2x + 3y = -11 \end{cases}$

9. $\begin{cases} 9x + y = 9 \\ 3x - 2y = -11 \end{cases}$



Math On the Spot
my.hrw.com

Solving Problems with Systems of Equations

Many real-world situations can be modeled with a system of equations.

EXAMPLE 3

Problem Solving



FL 8.EE.3.8c

The Simon family attended a concert and visited an art museum. Concert tickets were \$24.75 for adults and \$16.00 for children, for a total cost of \$138.25. Museum tickets were \$8.25 for adults and \$4.50 for children, for a total cost of \$42.75. How many adults and how many children are in the Simon family?



My Notes



Analyze Information

The answer is the number of adults and children.



Formulate a Plan

Solve a system to find the number of adults and children.



Solve

STEP 1

Choose variables and write a system of equations. Let x represent the number of adults. Let y represent the number of children.

Concert cost: $24.75x + 16.00y = 138.25$

Museum cost: $8.25x + 4.50y = 42.75$

STEP 2

Multiply both equations by 100 to eliminate the decimals.

$100(24.75x + 16.00y = 138.25) \rightarrow 2,475x + 1,600y = 13,825$

$100(8.25x + 4.50y = 42.75) \rightarrow 825x + 450y = 4,275$

© Houghton Mifflin Harcourt Publishing Company • Image Credits: © Education Images/UiG/Getty Images

STEP 3 Multiply the second equation by 3 and subtract this new equation from the first equation.

$$3(825x + 450y = 4,275)$$

Multiply each term in the second equation by 3 to get the same coefficients for the x-terms.

$$2,475x + 1,350y = 12,825$$

Simplify.

$$\begin{array}{r} 2,475x + 1,600y = 13,825 \\ -(2,475x + 1,350y = 12,825) \\ \hline \end{array}$$

Subtract the new equation from the first equation.

$$0x + 250y = 1,000$$

Subtract to eliminate the variable x.

$$250y = 1,000$$

Simplify and solve for y.

$$\frac{250y}{250} = \frac{1,000}{250}$$

Divide each side by 250.

$$y = 4$$

Simplify.

STEP 4 Substitute the solution into one of the original equations and solve for x.

$$8.25x + 4.50y = 42.75$$

Use the second equation.

$$8.25x + 4.50(4) = 42.75$$

Substitute 4 for the variable y.

$$8.25x + 18 = 42.75$$

Simplify.

$$8.25x = 24.75$$

Subtract 18 from each side.

$$x = 3$$

Divide each side by 8.25 and simplify.

STEP 5 Write the solution as an ordered pair: (3, 4).

There are 3 adults and 4 children in the family.



Justify and Evaluate

Substituting $x = 3$ and $y = 4$ into the original equations results in true statements. The answer is correct.

YOUR TURN

10. Contestants in the Run-and-Bike-a-thon run for a specified length of time, then bike for a specified length of time. Jason ran at an average speed of 5.2 mi/h and biked at an average speed of 20.6 mi/h, going a total of 14.2 miles. Seth ran at an average speed of 10.4 mi/h and biked at an average speed of 18.4 mi/h, going a total of 17 miles. For how long do contestants run and for how long do they bike?



Personal
Math Trainer

Online Assessment
and Intervention

my.hrw.com

Guided Practice

1. Solve the system $\begin{cases} 3x - y = 8 \\ -2x + 4y = -12 \end{cases}$ by multiplying and adding. (Example 1)

STEP 1 Multiply the first equation by 4. Add to the second equation.

$$4(3x - y = 8)$$

Multiply each term in the first equation by 4 to get opposite coefficients for the y -terms.

$$\boxed{}x - \boxed{}y = \boxed{}$$

Simplify.

$$+ \begin{array}{r} (-2x) + 4y = -12 \end{array}$$

Add the second equation to the new equation.

$$10x = \boxed{}$$

Add to eliminate the variable $\boxed{}$.

$$x = \boxed{}$$

Divide both sides by $\boxed{}$ and simplify.

STEP 2 Substitute into one of the original equations and solve for y .

$$y = \boxed{} \quad \text{So, } \boxed{} \text{ is the solution of the system.}$$

Solve each system of equations by multiplying first. (Examples 1, 2)

2. $\begin{cases} x + 4y = 2 \\ 2x + 5y = 7 \end{cases}$ _____

3. $\begin{cases} 3x + y = -1 \\ 2x + 3y = 18 \end{cases}$ _____

4. $\begin{cases} 2x + 8y = 21 \\ 6x - 4y = 14 \end{cases}$ _____

5. $\begin{cases} 2x + y = 3 \\ -x + 3y = -12 \end{cases}$ _____

6. $\begin{cases} 6x + 5y = 19 \\ 2x + 3y = 5 \end{cases}$ _____

7. $\begin{cases} 2x + 5y = 16 \\ -4x + 3y = 20 \end{cases}$ _____

8. Bryce spent \$5.26 on some apples priced at \$0.64 each and some pears priced at \$0.45 each. At another store he could have bought the same number of apples at \$0.32 each and the same number of pears at \$0.39 each, for a total cost of \$3.62. How many apples and how many pears did Bryce buy? (Example 3)

- a. Write equations to represent Bryce's expenditures at each store.

First store: _____ Second store: _____

- b. Solve the system.

Number of apples: _____ Number of pears: _____




ESSENTIAL QUESTION CHECK-IN

9. When solving a system by multiplying and then adding or subtracting, how do you decide whether to add or subtract?

8.4 Independent Practice



FL 8.EE.3.8b, 8.EE.3.8c



Personal Math Trainer

Online Assessment and Intervention

my.hrw.com

- 10. Explain the Error** Gwen used elimination with multiplication to solve the system
- $$\begin{cases} 2x + 6y = 3 \\ x - 3y = -1 \end{cases}$$
- Her work to find x is shown.
- Explain her error. Then solve the system.

$$\begin{aligned} 2(x - 3y) &= -1 \\ 2x - 6y &= -1 \\ +2x + 6y &= 3 \\ \hline 4x + 0y &= 2 \\ x &= \frac{1}{2} \end{aligned}$$

- 11. Represent Real-World Problems** At Raging River Sports, polyester-fill sleeping bags sell for \$79. Down-fill sleeping bags sell for \$149. In one week the store sold 14 sleeping bags for \$1,456.

Sleeping Bags

 <p>Nylon Down-filled, 35° \$149</p>	 <p>Flannel-lined Polyester-filled, 40° \$79</p>
--	--

- a.** Let x represent the number of polyester-fill bags sold and let y represent the number of down-fill bags sold. Write a system of equations you can solve to find the number of each type sold.

- b.** Explain how you can solve the system for y by multiplying and subtracting.

- c.** Explain how you can solve the system for y using substitution.

- d.** How many of each type of bag were sold?

- 12.** Twice a number plus twice a second number is 310. The difference between the numbers is 55. Find the numbers by writing and solving a system of equations. Explain how you solved the system.

- 13. Represent Real-World Problems** A farm stand sells apple pies and jars of applesauce. The table shows the number of apples needed to make a pie and a jar of applesauce. Yesterday, the farm picked 169 Granny Smith apples and 95 Red Delicious apples. How many pies and jars of applesauce can the farm make if every apple is used?

Type of apple	Granny Smith	Red Delicious
Needed for a pie	5	3
Needed for a jar of applesauce	4	2



FOCUS ON HIGHER ORDER THINKING

- 14. Make a Conjecture** Lena tried to solve a system of linear equations algebraically and in the process found the equation $5 = 9$. Lena thought something was wrong, so she graphed the equations and found that they were parallel lines. Explain what Lena's graph and equation could mean.

- 15.** Consider the system $\begin{cases} 2x + 3y = 6 \\ 3x + 7y = -1 \end{cases}$.

- a. Communicate Mathematical Ideas** Describe how to solve the system by multiplying the first equation by a constant and subtracting. Why would this method be less than ideal?

- b. Draw Conclusions** Is it possible to solve the system by multiplying both equations by integer constants? If so, explain how.

- c.** Use your answer from part b to solve the system.

Work Area

LESSON 8.5 Solving Special Systems

 **FL** 8.EE.3.8b

Solve systems of two linear equations in two variables algebraically, . . . Solve simple cases by inspection. Also 8.EE.3.8c



ESSENTIAL QUESTION

How do you solve systems with no solution or infinitely many solutions?

EXPLORE ACTIVITY

 **FL** 8.EE.3.8b

Solving Special Systems by Graphing

As with equations, some systems may have no solution or infinitely many solutions. One way to tell how many solutions a system has is by inspecting its graph.

Use the graph to solve each system of linear equations.

A $\begin{cases} x + y = 7 \\ 2x + 2y = 6 \end{cases}$


Is there a point of intersection? Explain.

Does this linear system have a solution? Use the graph to explain.

B $\begin{cases} 2x + 2y = 6 \\ x + y = 3 \end{cases}$

Is there a point of intersection? Explain.

Does this linear system have a solution? Use the graph to explain.



Reflect

- Use the graph to identify two lines that represent a linear system with exactly one solution. What are the equations of the lines? Explain your reasoning.

EXPLORE ACTIVITY (cont'd)

2. If each equation in a system of two linear equations is represented by a different line when graphed, what is the greatest number of solutions the system can have? Explain your reasoning.

3. Identify the three possible numbers of solutions for a system of linear equations. Explain when each type of solution occurs.



Solving Special Systems Algebraically

As with equations, if you solve a system of equations with no solution, you get a false statement, and if you solve a system with infinitely many solutions, you get a true statement.

EXAMPLE 1


FL 8.EE.3.8b

- A** Solve the system of linear equations by substitution.

$$\begin{cases} x - y = -2 \\ -x + y = 4 \end{cases}$$

STEP 1 Solve $x - y = -2$ for x :
 $x = y - 2$

STEP 2 Substitute the resulting expression into the other equation and solve.

$$-(y - 2) + y = 4$$


Substitute the expression for the variable x .

$$2 = 4$$

Simplify.

STEP 3 Interpret the solution. The result is the false statement $2 = 4$, which means there is no solution.

STEP 4 Graph the equations to check your answer. The graphs do not intersect, so there is no solution.



My Notes

B Solve the system of linear equations by elimination.


$$\begin{cases} 2x + y = -2 \\ 4x + 2y = -4 \end{cases}$$

STEP 1 Multiply the first equation by -2 .

$$-2(2x + y = -2) \rightarrow -4x + (-2y) = 4$$

STEP 2 Add the new equation from Step 1 to the original second equation.

$$\begin{array}{r} -4x + (-2y) = 4 \\ + \quad 4x + \quad 2y = -4 \\ \hline 0x + \quad 0y = 0 \\ 0 = 0 \end{array}$$



STEP 3 Interpret the solution. The result is the statement $0 = 0$, which is always true. This means that the system has infinitely many solutions.

STEP 4 Graph the equations to check your answer. The graphs are the same line, so there are infinitely many solutions.

Math Talk



Mathematical Practices

What solution do you get when you solve the system in part B by substitution? Does this result change the number of solutions? Explain.

Reflect

4. If x represents a variable and a and b represent constants so that $a \neq b$, interpret what each result means when solving a system of equations.

$x = a$ _____

$a = b$ _____

$a = a$ _____

5. In part B, can you tell without solving that the system has infinitely many solutions? If so, how?

YOUR TURN

Solve each system. Tell how many solutions each system has.

6. $\begin{cases} 4x - 6y = 9 \\ -2x + 3y = 4 \end{cases}$

7. $\begin{cases} x + 2y = 6 \\ 2x - 3y = 26 \end{cases}$

8. $\begin{cases} 12x - 8y = -4 \\ -3x + 2y = 1 \end{cases}$



Personal Math Trainer

Online Assessment and Intervention

my.hrw.com

Guided Practice

1. Use the graph to solve each system of linear equations. (Explore Activity)

A. $\begin{cases} 4x - 2y = -6 \\ 2x - y = 4 \end{cases}$

B. $\begin{cases} 4x - 2y = -6 \\ x + y = 6 \end{cases}$

C. $\begin{cases} 2x - y = 4 \\ 6x - 3y = 12 \end{cases}$


STEP 1

Decide if the graphs of the equations in each system intersect, are parallel, or are the same line.

System A: The graphs _____.

System B: The graphs _____.

System C: The graphs _____.



STEP 2

Decide how many points the graphs have in common.

Intersecting lines have _____ point(s) in common.

Parallel lines have _____ point(s) in common.

The same lines have _____ point(s) in common.

STEP 3

Solve each system.

System A has _____ points in common, so it has _____ solution.

System B has _____ point in common. That point is the solution, _____.

System C has _____ points in common. _____ ordered pairs on the line will make both equations true.

Solve each system. Tell how many solutions each system has. (Example 1)

2. $\begin{cases} x - 3y = 4 \\ -5x + 15y = -20 \end{cases}$

3. $\begin{cases} 6x + 2y = -4 \\ 3x + y = 4 \end{cases}$

4. $\begin{cases} 6x - 2y = -10 \\ 3x + 4y = -25 \end{cases}$



ESSENTIAL QUESTION CHECK-IN

5. When you solve a system of equations algebraically, how can you tell whether the system has zero, one, or an infinite number of solutions?

8.5 Independent Practice



FL 8.EE.3.8b, 8.EE.3.8c




Personal Math Trainer

Online Assessment and Intervention


my.hrw.com

Solve each system by graphing. Check your answer algebraically.

6.
$$\begin{cases} -2x + 6y = 12 \\ x - 3y = 3 \end{cases}$$



7.
$$\begin{cases} 15x + 5y = 5 \\ 3x + y = 1 \end{cases}$$



Solution: _____

Solution: _____

For Exs. 8–14, state the number of solutions for each system of linear equations.

8. a system whose graphs have the same slope but different y-intercepts

9. a system whose graphs have the same y-intercepts but different slopes

10. a system whose graphs have the same y-intercepts and the same slopes

11. a system whose graphs have different y-intercepts and different slopes

12. the system $\begin{cases} y = 2 \\ y = -3 \end{cases}$ _____

13. the system $\begin{cases} x = 2 \\ y = -3 \end{cases}$ _____

14. the system whose graphs were drawn using these tables of values:

Equation 1

x	0	1	2	3
y	1	3	5	7

Equation 2

x	0	1	2	3
y	3	5	7	9

15. **Draw Conclusions** The graph of a linear system appears in a textbook. You can see that the lines do not intersect on the graph, but also they do not appear to be parallel. Can you conclude that the system has no solution? Explain.

- 16. Represent Real-World Problems** Two school groups go to a roller skating rink. One group pays \$243 for 36 admissions and 21 skate rentals. The other group pays \$81 for 12 admissions and 7 skate rentals. Let x represent the cost of admission and let y represent the cost of a skate rental. Is there enough information to find values for x and y ? Explain.



- 17. Represent Real-World Problems** Juan and Tory are practicing for a track meet. They start their practice runs at the same point, but Tory starts 1 minute after Juan. Both run at a speed of 704 feet per minute. Does Tory catch up to Juan? Explain.



FOCUS ON HIGHER ORDER THINKING


- 18. Justify Reasoning** A linear system with no solution consists of the equation $y = 4x - 3$ and a second equation of the form $y = mx + b$. What can you say about the values of m and b ? Explain your reasoning.

- 19. Justify Reasoning** A linear system with infinitely many solutions consists of the equation $3x + 5 = 8$ and a second equation of the form $Ax + By = C$. What can you say about the values of A , B , and C ? Explain your reasoning.

- 20. Draw Conclusions** Both the points $(2, -2)$ and $(4, -4)$ are solutions of a system of linear equations. What conclusions can you make about the equations and their graphs?

Work Area


Ready to Go On?




8.1 Solving Systems of Linear Equations by Graphing

Solve each system by graphing.

1. $\begin{cases} y = x - 1 \\ y = 2x - 3 \end{cases}$



2. $\begin{cases} x + 2y = 1 \\ -x + y = 2 \end{cases}$



8.2 Solving Systems by Substitution

Solve each system of equations by substitution.

3. $\begin{cases} y = 2x \\ x + y = -9 \end{cases}$ _____

4. $\begin{cases} 3x - 2y = 11 \\ x + 2y = 9 \end{cases}$ _____

8.3 Solving Systems by Elimination

Solve each system of equations by adding or subtracting.

5. $\begin{cases} 3x + y = 9 \\ 2x + y = 5 \end{cases}$ _____

6. $\begin{cases} -x - 2y = 4 \\ 3x + 2y = 4 \end{cases}$ _____

8.4 Solving Systems by Elimination with Multiplication

Solve each system of equations by multiplying first.

7. $\begin{cases} x + 3y = -2 \\ 3x + 4y = -1 \end{cases}$ _____

8. $\begin{cases} 2x + 8y = 22 \\ 3x - 2y = 5 \end{cases}$ _____

8.5 Solving Special Systems

Solve each system. Tell how many solutions each system has.

9. $\begin{cases} -2x + 8y = 5 \\ x - 4y = -3 \end{cases}$ _____

10. $\begin{cases} 6x + 18y = -12 \\ x + 3y = -2 \end{cases}$ _____




ESSENTIAL QUESTION

11. What are the possible solutions to a system of linear equations, and what do they represent graphically?



Selected Response


1. The graph of which equation is shown?



- (A) $y = -2x + 2$ (C) $y = 2x + 2$
 (B) $y = -x + 2$ (D) $y = 2x + 1$
2. Which best describes the solutions to the system $\begin{cases} x + y = -4 \\ -2x - 2y = 0 \end{cases}$?
- (A) one solution (C) infinitely many
 (B) no solution (D) $(0, 0)$
3. Which of the following represents 0.000056023 written in scientific notation?
- (A) 5.6023×10^5 (C) 5.6023×10^{-4}
 (B) 5.6023×10^4 (D) 5.6023×10^{-5}
4. Which is the solution to $\begin{cases} 2x - y = 1 \\ 4x + y = 11 \end{cases}$?
- (A) $(2, 3)$ (C) $(-2, 3)$
 (B) $(3, 2)$ (D) $(3, -2)$

5. Which expression can you substitute in the indicated equation to solve $\begin{cases} 3x - y = 5 \\ x + 2y = 4 \end{cases}$?
- (A) $2y - 4$ for x in $3x - y = 5$
 (B) $4 - x$ for y in $3x - y = 5$
 (C) $3x - 5$ for y in $3x - y = 5$
 (D) $3x - 5$ for y in $x + 2y = 4$

6. What is the solution to the system of linear equations shown on the graph?



- (A) -1 (C) $(-1, -2)$
 (B) -2 (D) $(-2, -1)$
7. Which step could you use to start solving $\begin{cases} x - 6y = 8 \\ 2x - 5y = 3 \end{cases}$?
- (A) Add $2x - 5y = 3$ to $x - 6y = 8$.
 (B) Multiply $x - 6y = 8$ by 2 and add it to $2x - 5y = 3$.
 (C) Multiply $x - 6y = 8$ by 2 and subtract it from $2x - 5y = 3$.
 (D) Substitute $x = 6y - 8$ for x in $2x - 5y = 3$.

Mini-Task

8. A hot-air balloon begins rising from the ground at 4 meters per second at the same time a parachutist's chute opens at a height of 200 meters. The parachutist descends at 6 meters per second.
- a. Define the variables and write a system that represents the situation.
- _____
- _____
- b. Find the solution. What does it mean?
- _____
- _____

MODULE **7**

Solving Linear Equations

**ESSENTIAL QUESTION**

How can you use equations with variables on both sides to solve real-world problems?

EXAMPLE 1

A tutor gives students a choice of how to pay: a base rate of \$20 plus \$8 per hour, or a set rate of \$13 per hour. Find the number of hours of tutoring for which the cost is the same for either choice.

Plan 1 cost: $20 + 8x$ Plan 2 cost: $13x$

$$20 + 8x = 13x \quad \text{Write the equation.}$$

$$\begin{array}{r} 20 + 8x = 13x \\ -8x \quad -8x \\ \hline \end{array} \quad \text{Subtract } 8x \text{ from both sides.}$$

$$20 = 5x \quad \text{Divide both sides by 5.}$$

$$x = 4$$

The cost is the same for 4 hours of tutoring.

EXAMPLE 2

Solve $-2.4(3x + 5) = 0.8(x + 3.5)$.

$$-2.4(3x + 5) = 0.8(x + 3.5)$$

$$10(-2.4)(3x + 5) = 10(0.8)(x + 3.5) \quad \text{Multiply each side by 10 to clear some decimals.}$$

$$-24(3x + 5) = 8(x + 3.5)$$

$$-24(3x) - 24(5) = 8(x) + 8(3.5) \quad \text{Apply the Distributive Property.}$$

$$-72x - 120 = 8x + 28$$

$$\begin{array}{r} -72x - 120 = 8x + 28 \\ -8x \quad -8x \\ \hline \end{array} \quad \text{Subtract } 8x \text{ from both sides of the equation.}$$

$$-80x - 120 = 28$$

$$\begin{array}{r} -80x - 120 = 28 \\ +120 \quad +120 \\ \hline \end{array} \quad \text{Add 120 to both sides of the equation.}$$

$$-80x = 148$$

$$\begin{array}{r} -80x = 148 \\ -80 \quad -80 \\ \hline \end{array} \quad \text{Divide both sides of the equation by } -80.$$

$$x = -1.85$$

EXAMPLE 3

Solve $4(3x - 6) = 2(6x - 5)$.

$$4(3x - 6) = 2(6x - 5)$$

$$12x - 24 = 12x - 10$$

Apply the Distributive Property.

$$\underline{-12x} \quad \underline{-12x}$$

Subtract $8x$ from both sides of the equation.

$$-24 = -10$$

The statement is false.

There is no value of x that makes a true statement. Therefore, this equation has no solution.

EXERCISES

Solve. (Lessons 7.1, 7.2, 7.3, 7.4)

1. $13.02 - 6y = 8y$ _____

2. $\frac{1}{5}x + 5 = 19 - \frac{1}{2}x$ _____

3. $7.3t + 22 = 2.1t - 22.2$ _____

4. $1.4 + \frac{2}{5}e = \frac{3}{15}e - 0.8$ _____

5. $5(x - 4) = 2(x + 5)$ _____

6. $-7(3 + t) = 4(2t + 6)$ _____

7. $\frac{3}{4}(x + 8) = \frac{1}{3}(x + 27)$ _____

8. $3(4x - 8) = \frac{1}{5}(35x + 30)$ _____

9. $-1.6(2y + 15) = -1.2(2y - 10)$

10. $9(4a - 2) = 12(3a + 8)$

11. $6(x - \frac{1}{3}) = -2(x + 23)$

12. $8(p - 0.25) = 4(2p - 0.5)$

13. Write a real-world situation that could be modeled by the equation

$650 + 10m = 60m + 400$. (Lesson 7.1)

Solving Systems of Linear Equations



ESSENTIAL QUESTION

How can you use systems of equations to solve real-world problems?

Key Vocabulary

solution of a system of equations (*solución de un sistema de ecuaciones*)

system of equations (*sistema de ecuaciones*)

EXAMPLE 1 Solve the system of equations by substitution.

$$\begin{cases} 3x + y = 7 \\ x + y = 3 \end{cases}$$

Step 1 Solve an equation for one variable.

$$\begin{aligned} 3x + y &= 7 \\ y &= -3x + 7 \end{aligned}$$

Step 2 Substitute the expression for y in the other equation and solve.

$$\begin{aligned} x + y &= 3 \\ x + (-3x + 7) &= 3 \\ -2x + 7 &= 3 \\ -2x &= -4 \\ x &= 2 \end{aligned}$$

Step 3 Substitute the value of x into one of the equations and solve for the other variable, y .

$$\begin{aligned} x + y &= 3 \\ 2 + y &= 3 \\ y &= 1 \end{aligned}$$

(2, 1) is the solution of the system.

EXAMPLE 2 Solve the system of equations by elimination.

$$\begin{cases} x + y = 8 \\ 2x - 3y = 1 \end{cases}$$

Step 1 Multiply the first equation by 3 and add this new equation to the second equation.

$$\begin{aligned} 3(x + y = 8) &= 3x + 3y = 24 \\ 3x + 3y &= 24 \\ \underline{2x - 3y = 1} & \\ 5x + 0y &= 25 \\ 5x &= 25 \\ x &= 5 \end{aligned}$$

Step 2 Substitute the solution into one of the original equations and solve for y .

$$\begin{aligned} x + y &= 8 \\ 5 + y &= 8 \\ y &= 3 \end{aligned}$$

(5, 3) is the solution of the system.

EXERCISES

Solve each system of linear equations. (Lessons 8.1, 8.2, 8.3, 8.4, and 8.5)

14.
$$\begin{cases} x + y = -2 \\ 2x - y = 5 \end{cases}$$

15.
$$\begin{cases} y = 2x + 1 \\ x + 2y = 17 \end{cases}$$

16.
$$\begin{cases} y = -2x - 3 \\ 2x + y = 9 \end{cases}$$

17.
$$\begin{cases} y = 5 - x \\ 2x + 2y = 10 \end{cases}$$

18.
$$\begin{cases} 2x - y = 26 \\ 3x - 2y = 42 \end{cases}$$

19.
$$\begin{cases} 2x + 3y = 11 \\ 5x - 2y = 18 \end{cases}$$

20. Last week Andrew bought 3 pounds of zucchini and 2 pounds of tomatoes for \$7.05 at a farm stand. This week he bought 4 pounds of zucchini and 3 pounds of tomatoes, at the same prices, for \$9.83. What is the cost of 1 pound of zucchini and 1 pound of tomatoes at the farm stand?
-

Unit 3 Performance Tasks

1. **CAREERS IN MATH** **Hydraulic Engineer** A hydraulic engineer is studying the pressure in a particular fluid. The pressure is equal to the atmospheric pressure 101 kN/m plus 8 kN/m for every meter below the surface, where kN/m is kilonewtons per meter, a unit of pressure.
- a. Write an expression for the pressure at a depth of d_1 meters below the liquid surface. _____
- b. Write and solve an equation to find the depth at which the pressure is 200 kN/m. _____
- c. The hydraulic engineer alters the density of the fluid so that the pressure at depth d_2 below the surface is atmospheric pressure 101 kN/m plus 9 kN/m for every meter below the surface. Write an expression for the pressure at depth d_2 . _____
- d. If the pressure at depth d_1 in the first fluid is equal to the pressure at depth d_2 in the second fluid, what is the relationship between d_1 and d_2 ? Explain how you found your answer.
- _____
- _____
- _____



Selected Response


1. Ricardo and John start swimming from the same location. Ricardo starts 15 seconds before John and swims at a rate of 3 feet per second. John swims at a rate of 4 feet per second in the same direction as Ricardo. Which equation could you solve to find how long it will take John to catch up with Ricardo?

- (A) $4t + 3 = 3t$
- (B) $4t + 60 = 3t$
- (C) $3t + 3 = 4t$
- (D) $3t + 45 = 4t$

2. Gina and Rhonda work for different real estate agencies. Gina earns a monthly salary of \$5,000 plus a 6% commission on her sales. Rhonda earns a monthly salary of \$6,500 plus a 4% commission on her sales. How much must each sell to earn the same amount in a month?

- (A) \$1,500
- (B) \$15,000
- (C) \$75,000
- (D) \$750,000

3. What is the slope of the line?



- (A) -3
- (B) $-\frac{1}{3}$
- (C) $\frac{1}{3}$
- (D) 3

4. What is the solution of the system of equations?


$$\begin{cases} y = 2x - 3 \\ 5x + y = 11 \end{cases}$$

- (A) (2, 1)
- (B) (1, 2)
- (C) (3, -4)
- (D) (1, -1)

5. Alana is having a party. She bought 3 rolls of streamers and 2 packages of balloons for \$10.00. She realized she needed more supplies and went back to the store and bought 2 more rolls of streamers and 1 more package of balloons for \$6.25. How much did each roll of streamers and each package of balloons cost?

- (A) streamers: \$3.00, balloons: \$2.00
- (B) streamers: \$2.00, balloons: \$1.00
- (C) streamers: \$1.25, balloons: \$2.50
- (D) streamers: \$2.50, balloons: \$1.25

6. The triangle and the rectangle have the same perimeter.



Find the value of x .

- (A) 2
- (B) 10
- (C) 18
- (D) 24

7. What is the solution of the equation $8(3x + 4) = 2(12x - 8)$?
- (A) $x = -2$
 (B) $x = 2$
 (C) no solution
 (D) infinitely many solutions
8. A square wall tile has an area of 58,800 square millimeters. Between which two measurements is the length of one side?
- (A) between 24 and 25 millimeters
 (B) between 76 and 77 millimeters
 (C) between 242 and 243 millimeters
 (D) between 766 and 767 millimeters

Mini-Task

9. Lily and Alex went to a Mexican restaurant. Lily paid \$9 for 2 tacos and 3 enchiladas, and Alex paid \$12.50 for 3 tacos and 4 enchiladas.

- a. Write a system of equations that represents this situation.

- b. Use the system of equations to find how much the restaurant charges for a taco and for an enchilada.

- c. Describe the method you used to solve the system of equations.




Solutions of a system of two equations must make both equations true. Check solutions in both equations.

10. Use the system of equations to answer the questions below.

$$\begin{cases} 4x + 2y = -8 \\ 2x + y = 4 \end{cases}$$

- a. Graph the equations on the grid.



- b. How many solutions does the system of equations have? Explain your answer.

11. Isaac wants to join a gym. He checked out the membership fees at two gyms.

Gym A charges a new member fee of \$65 and \$20 per month.

Gym B charges a new member fee of \$25 and \$35 per month, but Isaac will get a discount of 20% on the monthly fee.

- a. Write an equation you can use to find the number of months for which the total costs at the gyms are the same.

- b. Solve the equation to find the number of months for which the total costs of the gyms are the same.
